

NDC 57896-317-06

GERI CARE®

ORIGINAL STRENGTH

# FAMOTIDINE TABLETS, USP 10 mg

## ACID REDUCER

**Just One Tablet**  
Prevents & Relieves Heartburn  
Due To Acid Indigestion

ACTUAL SIZE



Compare to active ingredient in  
Original Strength **PEPCID AC**®\*

**60**  
**TABLETS**

**Do not use if printed foil seal on bottle is broken or missing.**

### Drug Facts

**Active ingredient**      **Purpose**  
(in each tablet)  
Famotidine, USP 10 mg.....Acid reducer

#### Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

#### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.  
■ with other acid reducers

#### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

### Drug Facts (continued)

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

#### Other information

■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°C - 25°C (68°F - 77°F) ■ protect from moisture

NDC 57896-317-06

GERI CARE®

ORIGINAL STRENGTH

## FAMOTIDINE TABLETS USP 10 mg, ACID REDUCER

Distributed by: **Geri-Care Pharmaceuticals Corp.**  
1295 Towbin Avenue, Lakewood, NJ 08701  
Product of India      M.L. No.: 02/SKL/AP/2015/F/R

\*This product is not manufactured or distributed by the owner of the registered trademark **PEPCID AC**®, nor by the distributor of **PEPCID AC**®.      800229

### Drug Facts (continued)

#### Inactive ingredients

Microcrystalline Cellulose, Pre-gelatinized Starch, Sodium Starch Glycolate, Hydroxypropyl Cellulose, Magnesium Stearate, Hypromellose, Titanium Dioxide, Triacetin, Talc, Macrogol

#### Questions or comments?

1-800-540-3765 (toll-free)

**JUST ONE TABLET** prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages

### TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

REV 03/24

