Size: 3.75" x 6.875" **Colors: 4 Color Process**

Drug Facts

Active ingredient (per 10 mL=2 teaspoonfuls) Purposes Dextromethorphan HBr, USP 20 mg.... ..Cough suppressant Guaifenesin, USP 200 mg. .Expectorant

Uses • temporarily relieves cough due to minor throat and bronchial irritation • helps loosen phlegm (mucus) • helps thin bronchial secretions to make coughs more productive

Warnings

Do not use* if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

• if you have ever had an allergic reaction to any of the ingredients in this product.

Ask a doctor before use if you have
• a cough with too much phlegm (mucus) • a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

NDC 57896-291-16



GUAIFENESIN - Expectorant

- **CONTROLS COUGH**
- THINS & LOOSENS MUCUS
- RELIEVES CHEST CONGESTION

COMPARE TO THE ACTIVE INGREDIENTS OF ROBITUSSIN® DM*

SUGAR FREE • ALCOHOL FREE

COUGH SUPPRESSANT EXPECTORANT

For ages 12 years and over

16 FL OZ (473 mL)

Drug Facts (continued)

Directions

- take every 4 hours as needed, or as directed by a doctor do not take more than 6 doses in 24 hours
- do not exceed recommended dose

10 mL (2 teaspoonfuls) Adults and children 12 years and over Children under 12 years do not use

Other information

- each 10 mL contains: sodium 20 mg
 store at room temperature 15° 30°C (59°-86°F)
 protect from freezing do not refrigerate protect from light
 TAMPER-EVIDENT: Do not use if foil seal over bottle opening is torn, broken or missing.
- PHARMACIST: Preserve and dispense in a tight, light resistant container with a child resistant cap as defined in the USP.

Inactive ingredients: artificial and natural cherry flavor, citric acid, FD&C Red #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose.

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark Robitussin® DM. Each 10 mL dose of GERI-TUSSIN DM contains the same active ingredients as each 20 mL dose of Robitussin ® Cough + Chest Congestion DM.

For Institutional Use

Distributed by: GERI-CARE PHARMACEUTICALS CORP

1295 Towbin Avenue Lakewood, NJ 08701

Code# L-011 Lot#

Exp. Date:

REV::03/23