

IRON 65 mg
Plus Vitamin C
125 mg

DIETARY SUPPLEMENT

Compare to the dietary ingredients
in Vitron-C®*

60 TABLETS

Supplement Facts

Serving Size 1 Tablet

	Amount Per Serving	% Daily Value
Vitamin C (as ascorbic acid)	125 mg	139 %
IRON (as carbonyl iron)	65 mg	361 %

OTHER INGREDIENTS: Microcrystalline Cellulose, Croscarmellose Sodium, Magnesium Stearate, Hypromellose, PEG, Talc

Suggested use: Adults: take one (1) tablet daily or as recommended by your healthcare provider. Tablets should be swallowed whole. Not intended for use by children.

WARNING: Accidental overdose of iron containing products is a leading cause of fatal poisoning in children under 6. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.**

In case of accidental overdose, call a doctor or
Poison Control Center immediately.

WARNINGS: Do not exceed recommended dosage.

The treatment of any anemic condition should be under the advice and supervision of a physician. Since oral iron products interfere with the absorption of oral tetracycline antibiotics, these products should not be taken within two hours of each other. This product may occasionally cause constipation, diarrhea, or may color stool black. It may cause stomach upset (such as nausea), which may be minimized by taking with meals.

As with any supplement, if you are pregnant, nursing, taking medication, or have a medical condition, ask a health professional before use.
Store at room temperature.

TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

*This product is not manufactured or distributed by the owner of the registered trademark Vitron-C®.

DIST. BY: **GERI-CARE PHARMACEUTICALS CORP.**
1295 Towbin Avenue, Lakewood, NJ 08701

QUESTIONS OR COMMENTS? 1-800-540-3765

