

GERI-CARE®

NDC 57896-992-01

Indoor & Outdoor Allergies

CETIRIZINE HCl Tablets, USP

10 mg/Antihistamine

24-Hour Relief of:

- Sneezing
- Itchy, Watery Eyes
- Runny Nose
- Itchy Throat or Nose

Compares to Zyrtec® Allergy Original Prescription Strength*

100 TABLETS

10 mg each



DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient
(in each tablet)

Cetirizine HCl 10 mg.....Antihistamine

Purpose

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Uses

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- If breast-feeding: ask a health professional before use.
- If pregnant: not recommended.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Distributed by:
GERI-CARE PHARMACEUTICALS CORP.,
1295 TOWBIN AVE., LAKEWOOD, NJ 08701

* This product is not manufactured or distributed by Kenvue Inc., the owner of the registered trademark Zyrtec®.



3 57896 99201 9

RS3656 REV0325

Drug Facts (continued)

Directions

one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.	ask a doctor
Adults 65 years and over	ask a doctor
Children under 6 years of age	ask a doctor
Consumers with liver or kidney disease	ask a doctor

Other information

■ store between 20° to 25° C (68° to 77° F).

Inactive ingredients
hydrocellulose, lactose monohydrate, magnesium stearate, polyethylene glycol, polysorbate, povidone, pregelatinized starch, and titanium dioxide.

Questions or comments?
1-800-540-3765
Weekdays 9AM-5PM EST

Drug Facts (continued)

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- If breast-feeding: ask a health professional before use.
- If pregnant: not recommended.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.