NDC 57896-639-10



Aluminum Hydroxide, G Magnesium Hydroxide and Simethicone

GERI

Oral Suspension

1,200-1,200-120 mg/30 mL

100 x 30 mL Unit-Dose Cups

NDC 57896-639-10

Antacid & Antigas

1,200-1,200-120 mg/30 mL

Aluminum Hydroxide, Magnesium Hydroxide and Simethicone

Oral Suspension

Sugar Free

USUAL DOSAGE: See attached Drug Facts

FOR INSTITUTIONAL USE ONLY



100 x 30 mL Unit-Dose Cups

Purpose TAMPER EVIDENT: Do not use if lid is torn or broken. per 30 mL (1 Unit Dose) Active ingredients
Aluminum hydroxide 1,200 mg

Facts

Magnesium hydroxide 1,200 mg Simethicone 120 mg...... equivalent to dried gel USP)

Antacid Antigas

 heartburn • sour stomach • ac
 the symptoms referred to as gas relieves

acid indigestion

Warnings

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. a magnesium-restricted diet Ask a doctor before use if you have bidney disease • a magnesium-r

(eep out of reach of children. In case of overdose, get medical help or contact f pregnant or breast-feeding, ask a healthcare professional before use **Stop use and ask a doctor if** symptoms last for more than 2 weeks Poison Control Center right away (1-800-222-1222)

Antacids may interact with certain prescription drugs.

adults and children 12 years of age and older: take 30 mL, one unit dose, between do not take more than 4 unit dose (120 mL) in 24 hours or use the maximum dosage for more than 2 weeks children under 12 years; ask a doctor or as directed by a doctor

Other information

carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, store at room temperature benzyl alcohol, butylparaben, magnesium 510 mg, sodium 18 mg protect from freezing nactive ingredients

propy/paraben, purified water, saccharin sodium, sorbitol solution

Call 1-800-540-3765 comments? Questions

53/ Iss: 01/2024

LOT#

Exp. Date:

each 30 mL unit dose contains: