Drug Facts (continued)

Stop use and ask a doctor if
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lbs)	Under 24	24-35	36-47	48-59	60-71	72-95
Age (yr)	Under 2	2-3	4-5	6-8	9-10	11
Dose* (Tablets)	Ask a doctor	1	1 1/2	2	2 1/2	3

*or as directed by a doctor

Other information • store at 20-25°C (68-77°F); excursions permitted between 15°-30°C (59°-86°F) • avoid high humidity

Inactive ingredients: citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments? 1-800-540-3765

STOP PEELING

HEALTH STAR®

NDC 57896-155-24

CHILDREN'S CHEWABLE

ACETAMINOPHEN 160 mg

Ages 2-11

PAIN RELIEVER/ FEVER REDUCER

GRAPE FLAVOR

Compare to the active ingredient in TYLENOL® Children's Chewables*

24 Chewable Tablets

Chew or crush tablets completely before swallowing.

Drug Facts

TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

Active ingredient (in each chewable tablet)	Purposes
Acetaminophen 160 mg	Pain reliever / fever reducer

Uses • temporarily relieves minor aches and pains • temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

*This product is not manufactured or distributed by the owner of the registered trademark Tylenol®.

DIST. BY: **GERI-CARE** PHARMACEUTICALS CORP.
1295 Towbin Avenue, Lakewood, NJ 08701

REV 155-0726

3 57896 15524 7

PEEL HERE FOR MORE DRUG FACTS

Spot UV